



This is a repository copy of *Transethnic genome-wide association study provides insights in the genetic architecture and heritability of Long QT syndrome*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/161730/>

Version: Accepted Version

---

**Article:**

Lahrouchi, N, Tadros, R, Crotti, L et al. (103 more authors) (2020) Transethnic genome-wide association study provides insights in the genetic architecture and heritability of Long QT syndrome. *Circulation*. ISSN 0009-7322

<https://doi.org/10.1161/circulationaha.120.045956>

---

© 2020 American Heart Association, Inc. This is an author-produced version of a paper subsequently published in *Circulation*. Uploaded in accordance with the publisher's self-archiving policy.

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

# Transethnic genome-wide association study provides insights in the genetic architecture and heritability of long QT syndrome

Running title: Genome-wide association study in long QT syndrome

Najim Lahrouchi MD<sup>1-2\*</sup>, Rafik Tadros MD PhD<sup>1,3\*</sup>, Lia Crotti MD PhD<sup>2,4-7</sup>, Yuka Mizusawa MD PhD<sup>1-2</sup>, Pieter G. Postema<sup>1-2</sup> MD PhD, Leander Beekman BS<sup>1-2</sup>, Roddy Walsh PhD<sup>1-2</sup>, Kanae Hasegawa MD PhD<sup>8-9</sup>, Julien Barc PhD<sup>2,10</sup>, Marko Ernstring PhD<sup>2,11</sup>, Kari L. Turkowski BS<sup>12</sup>, Andrea Mazzanti MD PhD<sup>2,13</sup>, Britt M. Beckmann MD<sup>14</sup>, Keiko Shimamoto MD<sup>15</sup>, Ulla-Britt Diamant MD PhD<sup>2,16</sup>, Yanushi D. Wijeyeratne MD<sup>2,17</sup>, Yu Kucho MD<sup>18</sup>, Tomas Robyns MD PhD<sup>2,19-20</sup>, Taisuke Ishikawa PhD<sup>21</sup>, Elena Arbelo MD PhD<sup>22</sup>, Michael Christiansen MD<sup>23-25</sup>, Annika Winbo MD PhD<sup>26</sup>, Reza Jabbari MD PhD<sup>2,27</sup>, Steven A. Lubitz MD PhD<sup>28-29</sup>, Johannes Steinfurt MD<sup>30</sup>, Boris Rudic MD<sup>31</sup>, Bart Loeys MD PhD<sup>32</sup>, M. Ben Shoemaker MD<sup>33</sup>, Peter E. Weeke MD PhD<sup>27,33</sup>, Ryan Pfeiffer BS<sup>34</sup>, Brianna Davies MS<sup>35</sup>, Antoine Andorin MD<sup>17,36</sup>, Nynke Hofman PhD<sup>1-2</sup>, Federica Dagradi MD<sup>2,4</sup>, Matteo Pedrazzini PhD<sup>5</sup>, David J. Tester BS<sup>12</sup>, Johan M. Bos MD PhD<sup>12</sup>, Georgia Sarquella-Brugada MD PhD<sup>2,37-39</sup>, Óscar Campuzano PhD<sup>39-41</sup>, Pyotr G. Platonov MD PhD<sup>42</sup>, Birgit Stallmeyer MD<sup>11</sup>, Sven Zumhagen MD<sup>11</sup>, Eline A. Nannenbergh MD PhD<sup>43</sup>, Jan H. Veldink MD PhD<sup>44</sup>, Leonard H. van den Berg MD PhD<sup>44</sup>, Ammar Al-Chalabi MD PhD<sup>45-46</sup>, Christopher E. Shaw MD PhD<sup>46-47</sup>, Pamela J. Shaw MD<sup>48</sup>, Karen E. Morrison MD PhD<sup>49</sup>, Peter M. Andersen MD<sup>50</sup>, Martina Müller-Nurasyid PhD<sup>14,51</sup>, Daniele Cusi PhD<sup>52-53</sup>, Cristina Barlassina PhD<sup>52-53</sup>, Pilar Galan MD PhD<sup>54</sup>, Mark Lathrop PhD<sup>55</sup>, Markus Munter PhD<sup>55</sup>, Thomas Werge PhD<sup>56-58</sup>, Marta Ribasés PhD<sup>59</sup>, Tin Aung MD PhD<sup>60</sup>, Chiea C. Khor PhD<sup>61</sup>, Mineo Ozaki MD PhD<sup>62</sup>, Peter Lichtner PhD<sup>63</sup>, Thomas Meitinger MD<sup>63</sup>, J. Peter van Tintelen MD PhD<sup>43,64,65</sup>, Yvonne Hoedemaekers MD PhD<sup>64</sup>, Isabelle Denjoy MD<sup>2,66</sup>, Antoine Leenhardt MD<sup>2,66</sup>, Carlo Napolitano MD PhD<sup>2,13</sup>, Wataru Shimizu MD PhD<sup>15,67</sup>, Jean-Jacques Schott PhD<sup>2,10,36</sup>, Jean-Baptiste Gourraud MD PhD<sup>2,10,36</sup>, Takeru Makiyama MD PhD<sup>68</sup>, Seiko Ohno MD PhD<sup>8,69-70</sup>, Hideki Itoh<sup>8,69</sup>, Andrew D. Krahn MD<sup>35</sup>, Charles Antzelevitch PhD<sup>71-73</sup>, Dan M. Roden MD PhD<sup>74</sup>, Johan Saenen MD PhD<sup>75</sup>, Martin Borggrefe MD<sup>31</sup>, Katja E. Odening MD<sup>30</sup>, Patrick T. Ellinor MD PhD<sup>28,29</sup>, Jacob Tfelt-Hansen MD<sup>2,27,76</sup>, Jonathan R. Skinner MD<sup>77</sup>, Maarten P. van den Berg MD PhD<sup>78</sup>, Morten Salling Olesen PhD<sup>79-80</sup>, Josep Brugada MD PhD<sup>81</sup>, Ramón Brugada MD PhD<sup>40,82-83</sup>, Naomasa Makita MD PhD<sup>84</sup>, Jeroen Breckpot MD PhD<sup>2,85</sup>, Masao Yoshinaga MD PhD<sup>18</sup>, Elijah R. Behr MD<sup>2,17</sup>, Annika Rydberg MD PhD<sup>2,16</sup>, Takeshi Aiba MD PhD<sup>15</sup>, Stefan Käb MD PhD<sup>14</sup>, Silvia G. Priori MD PhD<sup>2,13</sup>, Pascale Guicheney PhD<sup>86</sup>, Hanno L. Tan MD PhD<sup>1,2,87</sup>, Christopher Newton-Cheh MD<sup>88</sup>, Michael J. Ackerman MD PhD<sup>12</sup>, Peter J. Schwartz MD<sup>2,4-5</sup>, Eric Schulze-Bahr MD<sup>2,11</sup>, Vincent Probst MD PhD<sup>2,10,67</sup>, Minoru Horie MD PhD<sup>8,69</sup>, Arthur A. Wilde MD PhD<sup>1,2</sup>, Michael W.T. Tanck PhD<sup>89</sup>, Connie R. Bezzina PhD<sup>1,2</sup> \*These authors equally contributed to the study and share first authorship

## Corresponding author:

Connie R. Bezzina

Amsterdam UMC, AMC Heart Center, Dept. Clin. & Exp. Cardiology

Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands

[c.r.bezzina@amc.uva.nl](mailto:c.r.bezzina@amc.uva.nl)

## Affiliations

1. Amsterdam UMC, University of Amsterdam, Heart Center; Department of Clinical and Experimental Cardiology, Amsterdam Cardiovascular Sciences, Meibergdreef 9, Amsterdam, The Netherlands
2. Member of the European Reference Network for rare, low prevalence and complex diseases of the heart - ERN GUARD-Heart
3. Cardiovascular Genetics Center, Montreal Heart Institute and Faculty of Medicine, Université de Montréal, Montreal, Canada
4. Istituto Auxologico Italiano, IRCCS, Center for Cardiac Arrhythmias of Genetic Origin, Milan, Italy
5. Istituto Auxologico Italiano, IRCCS, Laboratory of Cardiovascular Genetics, Milan, Italy
6. Istituto Auxologico Italiano, IRCCS, Department of Cardiovascular, Neural and Metabolic Sciences, San Luca Hospital, Milan, Italy
7. Department of Medicine and Surgery, University of Milano-Bicocca, Milan, Italy
8. Department of Cardiovascular Medicine, Shiga University of Medical Science, Otsu, Japan
9. Department of Cardiovascular Medicine, Faculty of Medical Sciences, University of Fukui, Fukui, Japan
10. l'institut du thorax, INSERM, CNRS, UNIV Nantes, Nantes, France
11. Institute for Genetics of Heart Diseases, Department of Cardiovascular Medicine, University Hospital Muenster, Muenster, Germany
12. Departments of Cardiovascular Medicine (Division of Heart Rhythm Services and the Windland Smith Rice Genetic Heart Rhythm Clinic), Pediatric and Adolescent Medicine (Division of Pediatric Cardiology), and Molecular Pharmacology & Experimental Therapeutics (Windland Smith Rice Sudden Death Genomics Laboratory), Mayo Clinic, Rochester, MN, USA
13. Molecular Cardiology, ICS Maugeri, IRCCS and Department of Molecular Medicine, University of Pavia, Pavia, Italy
14. Department of Internal Medicine I, University Hospital of the Ludwig Maximilians University, Munich, Germany
15. Department of Cardiovascular Medicine, National Cerebral and Cardiovascular Center, Osaka, Japan
16. Department of Clinical Sciences, Unit of Paediatrics, Umeå University, Umeå, Sweden
17. Molecular and Clinical Sciences Research Institute, St George's University of London, London, United Kingdom; Cardiology Clinical Academic Group, St George's University Hospitals NHS Foundation Trust, London, United Kingdom
18. National Hospital Organization Kagoshima Medical Center, Japan
19. Department of cardiovascular diseases, University Hospitals Leuven, Belgium
20. Department of cardiovascular sciences, KU Leuven, Belgium
21. Omics Research Center, National Cerebral and Cardiovascular Center, Osaka, Japan
22. Cardiovascular Institute, Hospital Clinic de Barcelona, Universitat de Barcelona, Spain. Institut d'Investigació August Pi i Sunyer (IDIBAPS). Centro de Investigación Biomedica en Red de Enfermedades Cardiovasculares (CIBERCV), Barcelona, Spain
23. Department of Congenital Disorders, Statens Serum Institute, Copenhagen, Denmark
24. The Lundbeck Foundation Initiative for Integrative Psychiatric Research, iPSYCH, Copenhagen, Denmark
25. Laboratory of Experimental Cardiology, Department of Biomedical Sciences, University of Copenhagen, Copenhagen, Denmark
26. Department of Physiology, The University of Auckland, Auckland, New Zealand
27. The Department of Cardiology, The Heart Centre, Copenhagen University Hospital, Rigshospitalet, Copenhagen, Denmark.
28. Cardiac Arrhythmia Service, Massachusetts General Hospital, Boston, Massachusetts, USA
29. Cardiovascular Disease Initiative and Program in Medical and Population Genetics, Broad Institute, Cambridge, Boston, Massachusetts, USA

30. Department of Cardiology and Angiology I, Heart Center University of Freiburg, Medical Faculty, Freiburg, Germany
31. Department of Medicine, University Medical Center Mannheim, Mannheim, Germany; German Center for Cardiovascular Research (DZHK), Partner Site Heidelberg/Mannheim
32. Department of Clinical Genetics, Antwerp University Hospital, Antwerp, Belgium
33. Department of Medicine, Vanderbilt University Medical Center, Nashville, Tennessee.
34. Masonic Medical Research Institute, Utica, NY, USA
35. Heart Rhythm Services, Division of Cardiology, Department of Medicine, University of British Columbia, Vancouver, British Columbia, Canada
36. l'institut du thorax, CHU Nantes, Service de Cardiologie, Nantes, France
37. Arrhythmia, Inherited Heart Disease and Sudden Death Unit, Hospital Sant Joan de Déu, European Reference Center at the ERN GUARD-Heart Reference Network for Rare Cardiac Diseases, Barcelona, Spain
38. Medical Science Department, School of Medicine, University of Girona, Girona, Spain
39. Cardiovascular Program, Research Institute of Sant Joan de Déu (IRSJD), Barcelona, Spain
40. Center for Biomedical Diagnosis, Hospital Clinic de Barcelona, Universitat de Barcelona, Spain. Institut d'Investigació August Pi i Sunyer (IDIBAPS). Cardiovascular Genetics Center, University of Girona-IDIBGI, Girona (Spain). Medical Science Department, School of Medicine, University of Girona, Girona, Spain
41. Centro de Investigación Biomédica en Red de Enfermedades Cardiovasculares (CIBERCV), Madrid, Spain
42. Center for Integrative Electrophysiology (CIEL), Department of Cardiology, Clinical Sciences, Lund University, Lund, Sweden
43. Department of Clinical Genetics, Amsterdam UMC, University of Amsterdam, Amsterdam, the Netherlands
44. Department of Neurology, UMC Utrecht Brain Center, University Medical Center Utrecht, Utrecht, The Netherlands
45. King's College Hospital, Bessemer Road, London, SE5 9RS, UK
46. Department of Basic and Clinical Neuroscience, King's College London, Maurice Wohl Clinical Neuroscience Institute, London, UK
47. UK Dementia Research Institute, King's College London, London, UK
48. Sheffield Institute for Translational Neuroscience, University of Sheffield, Sheffield, UK
49. Faculty of Medicine, University of Southampton, University Hospital Southampton, Southampton, UK
50. Department of Neurology, Ulm University, 89081 Ulm, Germany; and Department of Pharmacology and Clinical Neuroscience, Umeå University, 90187 Umeå, Sweden
51. Institute of Genetic Epidemiology, Helmholtz Zentrum München–German Research Center for Environmental Health, Neuherberg, Germany; Chair of Genetic Epidemiology, IBE, Faculty of Medicine, LMU Munich, Germany.
52. Department of Health Sciences, University of Milan, Milan, Italy
53. Bio4Dreams - business nursery for life sciences, Milan, Italy
54. Equipe de Recherche en Epidémiologie Nutritionnelle, Centre d'Epidémiologie et Statistiques Paris Cité, Université Paris 13, Inserm (U1153), Inra (U1125), COMUE Sorbonne-Paris-Cité, Bobigny, France
55. McGill University and Génome Québec Innovation Centre, Montréal, Québec, Canada
56. The Lundbeck Foundation Initiative for Integrative Psychiatric Research, iPSYCH, Copenhagen, Denmark
57. Institute of Biological Psychiatry, Mental Health Centre Sct Hans, Copenhagen University Hospital, Roskilde, Denmark
58. Department of Clinical Medicine, University of Copenhagen, Copenhagen, Denmark.
59. Psychiatric Genetics Unit, Institut Vall d'Hebron Research (VHIR), Universitat Autònoma de Barcelona, Barcelona, Spain

60. Singapore Eye Research Institute, Singapore, Singapore
61. Genome Institute of Singapore, Singapore, Singapore
62. Ozaki Eye Hospital, 1-15, Kamezaki, Hyuga, Miyazaki 883-0066 Japan
63. Institute of Human Genetics, Helmholtz Zentrum München, Neuherberg, Germany.
64. Department of Clinical Genetics, University Medical Centre Groningen, Groningen, The Netherlands
65. Department of Clinical Genetics, University Medical Centre Utrecht, University of Utrecht, Utrecht, The Netherlands
66. AP-HP, Hôpital Bichat, Département de Cardiologie et Centre de Référence des Maladies Cardiaques Héréditaires, F-75018 Paris, France, Université de Paris INSERM U1166, F-75013 Paris, France
67. Department of Cardiovascular Medicine, Graduate School of Medicine, Nippon Medical School, Tokyo, Japan
68. Department of Cardiovascular Medicine, Kyoto University Graduate School of Medicine, Kyoto, Japan
69. Center for Epidemiologic Research in Asia, Shiga University of Medical Science, Otsu, Japan
70. Department of Bioscience and Genetics, National Cerebral and Cardiovascular Center, Suita, Japan
71. Lankenau Institute for Medical Research, Wynnewood, PA, USA
72. Lankenau Heart Institute, Wynnewood, PA, USA
73. Sidney Kimmel Medical College, Thomas Jefferson University, Philadelphia, PA, USA
74. Department of Biomedical Informatics, Vanderbilt University Medical Center, Nashville, TN, USA; Department of Medicine, Vanderbilt University Medical Center, Nashville, TN, USA; Department of Pharmacology, Vanderbilt University Medical Center, Nashville, TN, USA.
75. Department of Cardiology, Antwerp University Hospital, Belgium
76. Department of Forensic Medicine, Faculty of Medical Sciences, University of Copenhagen, Copenhagen, Denmark
77. Cardiac Inherited Disease Group, Starship Childrens Hospital, Auckland, New Zealand
78. Department of cardiology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands
79. Laboratory for Molecular Cardiology, Department of Cardiology, The Heart Centre, Rigshospitalet (Copenhagen University Hospital), Copenhagen, Denmark
80. Department of Biomedical Sciences, University of Copenhagen, Copenhagen, Denmark
81. Arrhythmia Unit, Hospital Sant Joan de Déu, University of Barcelona, Barcelona (Spain). Cardiovascular Institute, Hospital Clinic de Barcelona, Universitat de Barcelona, Spain. Institut d'Investigació August Pi i Sunyer (IDIBAPS).
82. Cardiovascular Genetics Center, University of Girona-IDIBGI, Girona, Spain. Medical Science Department, School of Medicine, University of Girona, Girona, Spain
83. Cardiology Service, Hospital Josep Trueta, Girona, Spain
84. National Cerebral and Cardiovascular Center Research Institute, Osaka, Japan
85. Centre for human genetics, university hospitals Leuven, Belgium
86. INSERM, Sorbonne University, UMRS 1166, Institute of Cardiometabolism and Nutrition (ICAN), Paris, France
87. Netherlands Heart Institute, Utrecht, The Netherlands
88. Cardiovascular Research Center and Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA, USA
89. Department of Clinical Epidemiology, Biostatistics and Bioinformatics, Amsterdam UMC, University of Amsterdam. Amsterdam, The Netherlands

## ABSTRACT

**Background:** Long QT syndrome (LQTS) is a rare genetic disorder and a major preventable cause of sudden cardiac death in the young. A causal rare genetic variant with large effect size is identified in up to 80% of probands (genotype positive) and cascade family screening shows incomplete penetrance of genetic variants. Furthermore, a proportion of cases meeting diagnostic criteria for LQTS remain genetically elusive despite genetic testing of established genes (genotype negative). These observations raise the possibility that common genetic variants with small effect size contribute to the clinical picture of LQTS. This study aimed to characterize and quantify the contribution of common genetic variation to LQTS disease susceptibility.

**Methods:** We conducted genome-wide association studies (GWAS) followed by transethnic meta-analysis in 1,656 unrelated LQTS patients of European or Japanese ancestry and 9,890 controls to identify susceptibility single nucleotide polymorphisms (SNPs). We estimated the SNP heritability ( $h^2_{\text{SNP}}$ ) of LQTS and tested the genetic correlation between LQTS susceptibility and other cardiac traits. Furthermore, we tested the aggregate effect of the 68 SNPs previously associated with QTc in the general population using a polygenic risk score (PRS<sub>QT</sub>).

**Results:** Genome-wide association analysis identified three loci associated with LQTS at genome-wide statistical significance ( $P < 5 \times 10^{-8}$ ) near *NOS1AP*, *KCNQ1* and *KLF12*, and one missense variant in *KCNE1* (p.Asp85Asn) at the suggestive threshold ( $P < 10^{-6}$ ). Heritability analyses showed that ~15% of variance in overall LQTS susceptibility was attributable to common genetic variation ( $h^2_{\text{SNP}}$  0.148; standard error [SE] 0.019). LQTS susceptibility showed a strong genome-wide genetic correlation with the QT interval in the general population ( $r_g = 0.40$ ,  $P = 3.2 \times 10^{-3}$ ). PRS<sub>QT</sub> was greater in LQTS cases compared to controls ( $P < 10^{-13}$ ), and notably, among LQTS patients PRS<sub>QT</sub> was greater in genotype negative compared to genotype positive patients ( $P < 0.005$ ).

**Conclusion:** This work establishes an important role for common genetic variation in susceptibility to LQTS. We demonstrate overlap between genetic control of the QT interval in the general population and genetic factors contributing to LQTS susceptibility. Using polygenic risk score analyses aggregating common genetic variants that modulate the QT interval in the general population, we provide evidence for a polygenic architecture in genotype negative LQTS.

**Key words:** long QT syndrome, genome-wide association study, polygenic risk score, heritability, QT-interval

## **Non-standard Abbreviations and Acronyms**

ACMG - American College of Medical Genetics and Genomics

AMP - Association of Molecular Pathology

GWAS - Genome-wide association study

HRC - Haplotype reference consortium

gnomAD - Genome Aggregation Database

LAE - Life-threatening arrhythmic event

LQTS - Long QT syndrome

MAF - Minor allele frequency

PRS - Polygenic risk score

QC - Quality control

SCD - Sudden cardiac death

SNP - Single nucleotide polymorphisms

VUS - Variant of unknown significance



**What is new?**

- A genome-wide association study in Long-QT syndrome (LQTS) patients establishes and quantifies the role of common genetic variation in susceptibility to LQTS.
- Genetic overlap exists between control of QT-interval in the general population and susceptibility to LQTS.
- Polygenic risk score analyses based on common genetic variants that modulate the QT-interval in the general population provide evidence for a polygenic architecture in LQTS patients that remain genetically elusive despite genetic testing of established genes (i.e. genotype negative).

**What are the clinical implications?**

- These findings enhance the understanding of the genetic basis of LQTS and underscore the genetic relationship between QT-interval in the general population and susceptibility to LQTS.
- Increasing burden of QT-prolonging common variants is associated with higher susceptibility for LQTS.
- Polygenicity in genotype negative LQTS patients implies that risk is not primarily attributable to one genetic factor inherited from one of the biological parents as is the case for autosomal dominant LQTS.
- Future clinical utility of genetic testing based on polygenic inheritance necessitates the availability of polygenic risk scores with high discriminative capacity.

## INTRODUCTION

Long QT syndrome (LQTS) is a rare inherited disorder of ventricular repolarization characterized by prolongation of the QT interval on the electrocardiogram (ECG).[1,2] LQTS has a prevalence of approximately 1 in 2500, and is a major and often preventable cause of sudden cardiac death (SCD) in the young.[3,4] Multiple genes have been implicated in LQTS and clinical genetic testing is now performed to identify causative rare genetic variants.[5] Disease-causing variants (i.e. mutations) in the three major LQTS genes, i.e. *KCNQ1* (LQT1), *KCNH2* (LQT2) and *SCN5A* (LQT3), account for up to 80% of LQTS cases overall and >95% of genotype positive LQTS.[2]

Studies in families with multiple mutation carriers have shown that disease penetrance (proportion of carriers that manifest with a prolonged QT interval) can be low,[6–8] and that among those with disease manifestations, there can be broad variability in the types of symptoms and severity thereof (variable expression).[2,6–8] These observations suggest that, like other Mendelian disorders, allocating the disease in the individual patient exclusively to a rare variant at a single locus (i.e. monogenetic) might be an oversimplification of biological phenomena. It is likely that a combination of genetic and non-genetic modifying factors underlies this clinical variability. A comprehensive knowledge of such risk factors that affect penetrance and expressivity of disease-causing variants in LQTS will improve the predictive accuracy of genetic testing in the individual patient and enable personalized clinical interventions. While many clinical risk factors such as sex, hypokalaemia, or bradyarrhythmia, have been implicated as modulators of the clinical manifestations of LQTS[9], modulatory genetic factors remain largely unexplored with the exception of a few proof-of-concept studies using a candidate gene approach.[10–14]

Besides variability in disease manifestations among carriers of pathogenic variants, an outstanding issue in LQTS is the fact that in ~20% of patients, an underlying causal rare genetic variant remains unidentified after extensive panel-based genetic testing.[15] This complicates cascade screening in families and the pre-symptomatic identification of affected relatives.

Although a small proportion of such genotype negative LQTS patients could have a yet unknown Mendelian defect, another possibility is that a more complex inheritance pattern underlies the disorder in a subset of these patients.

Previous work has shown that genome-wide association study (GWAS) comparing cases of a rare arrhythmia syndrome to unaffected controls can define modulators of disease susceptibility and suggest a polygenic etiology.[16] We report here a GWAS in ~1,700 unrelated LQTS patients of European or Japanese ancestry, identifying common genetic variants implicated in LQTS disease susceptibility, and providing a quantification of the contribution of common genetic variants to LQTS predisposition. Using polygenic risk score analyses aggregating common genetic variants that modulate the QT interval in the general population, we provide evidence for a polygenic architecture in genotype negative LQTS.

## METHODS

The summary statistics generated in this study are available from the corresponding author upon request.

### Study population

We established an international consortium allowing recruitment of 1,781 unrelated LQTS patients: 1,344 cases of European ancestry from 23 referral centres in Europe, New Zealand and North America, as well as 437 patients of East Asian ancestry from 4 referral centres in Japan (**Table I in the Supplement**). Included unrelated individuals were probands (97%) except when DNA was not available, in which case one other affected family member was included instead. Included LQTS patients had a clinical diagnosis of LQTS[5] and were classified as “genotype positive” if they carried a single rare variant in one of the three established major LQTS genes [*KCNQ1* (LQT1), *KCNH2* (LQT2) and *SCN5A* (LQT3)], or “genotype negative” if no rare variant was identified in genes unequivocally associated with non-syndromic LQTS (*KCNQ1*, *KCNH2*, *SCN5A*, *CALM1-3* and *TRDN*).[17–19] A rare variant was defined as a protein sequence altering (i.e. missense, nonsense, frameshift deletion, in-frame deletion, large deletion and duplication) or splice-site variant with an allele frequency  $<1 \times 10^{-4}$  in the Genome Aggregation Database (gnomAD).[20–22] Genetic testing and variant curation as per the American College of Medical Genetics and Genomics and Association of Molecular Pathology (ACMG/AMP) guidelines[23] was conducted as in the **Supplementary Methods**. All subjects or their guardians provided informed consent, and the study was approved by the appropriate ethical review boards.

### Phenotypic characterization and measurement of the QT interval

Clinical data were collected at each of the participating centres. We collected a baseline ECG for each patient, preferably not during beta-blocker use. The QT-interval duration was measured as previously described (**Figure I in the Supplement, Supplementary**

**Methods**).[24] In genotype negative patients, a LQTS diagnosis was additionally curated by 2 clinicians (NL, RT) and in case of uncertainty, two senior LQTS experts (AAW, PJS) were consulted. As per international guidelines[5], we only included genotype negative patients with a LQTS risk score  $\geq 3.5$  or with a resting QTc  $\geq 500$ ms in repeated 12-lead ECGs, in the absence of a secondary cause for QT prolongation.

### **Genome-wide array genotyping, quality control and imputation**

We performed genome-wide genotyping for all European cases on the Illumina HumanOmniExpress array and for all Japanese cases on the Illumina Global Screening Array. Genotypic data of 8,219 control individuals of European ancestry and 1,671 individuals of Japanese ancestry were obtained from different cohorts (**Table II in the Supplement**). Quality control (QC)[25], imputation and association analysis were performed separately in the European and Japanese datasets. All genetic variants were mapped to and reported using Genome Reference Consortium Human genome build 37 (GRCh37).

After QC (see **Supplementary Methods** for details), we performed genome-wide imputation using Eagle2 phasing, Minimac3 and the Haplotype reference consortium (HRCr1.1) panel implemented on the Michigan Imputation Server for both the European and Japanese datasets.[26] After imputation, only single nucleotide polymorphisms (SNPs) with MAF>0.01 and a Minimac3 imputation score of  $R^2 > 0.3$  were included in further analyses.

### **Genome-wide association analysis**

We performed genome-wide association analyses to assess the role of common variants in LQTS susceptibility (case-control) and severity (QTc within the cases). Case-control association of alternate allele dosage with LQTS was performed using logistic regression correcting for genotypic PC 1-10. Quantitative trait analyses for QTc were conducted using a linear regression model correcting for age, beta-blocker use at ECG, LQTS type ([*KCNQ1* (*LQT1*), *KCNH2* (*LQT2*) and *SCN5A* (*LQT3*), or genotype negative), sex, and PC 1-10. Genome-wide association analyses were carried out separately for the European and

Japanese LQTS cohorts, followed by meta-analysis using an inverse variance weighted fixed effect model, implemented in METAL (version 2011-03-25).[27] Genome-wide statistical significance and suggestive thresholds were set to  $P < 5 \times 10^{-8}$  and  $P < 1 \times 10^{-6}$ , respectively. Summary statistics were uploaded to FUMA (Functional Mapping and Annotation of GWAS) for generation of Manhattan, quantile-quantile and regional association plots for risk loci.[28]

### **Survival analyses**

Time to life-threatening arrhythmic events (LAE) survival analyses were performed in the LQTS cases. Follow up started at birth and stopped at the date of a document LAE, the last visit or the 41<sup>st</sup> birthday, whichever came first. LAE were defined as out of hospital cardiac arrest (OHCA) or hemodynamically unstable ventricular tachycardia/ventricular fibrillation (VT/VF) or appropriate implantable cardioverter-defibrillator (ICD) therapy. The effect of genotype positive versus genotype negative status was estimated using Cox proportional hazards regression with/without adjustment for classic risk factors, i.e. sex and  $QTc \geq 500$  ms. To examine possible differences in effect of these well-recognized risk factors in genotype positive and genotype negative LQTS cases, interactions between these risk factors and genotype status were included in the model. In addition, puberty and a sex  $\times$  puberty interaction were included to model the modifying effect of puberty on the effect of sex. Puberty was included as time-varying covariate and the age of puberty was set at 16 years in both sexes (i.e. during the follow up period prior to the age of 16, puberty was coded as 0, whereas puberty was coded as 1 during the remainder of the follow up period). Kaplan Meier curves were created to illustrate the cumulative event free survival and log rank tests were used to compare the survival curves.

### **Polygenic risk scores (PRS)**

For all cases and controls, we calculated a weighted QT polygenic risk score ( $PRS_{QT}$ ) comprising 68 SNPs that had been associated with the QT-interval in the general population at genome-wide statistical significance, in a study primarily including Europeans.[29] All 68 SNPs were included in the European dataset analyses whereas only 60/68 SNPs were well-

imputed and included in the Japanese dataset analyses (**Table III in the Supplement**).  $PRS_{QT}$  was calculated by multiplying the alternate allele dosage by the associated effect size ( $\beta$ ) in the published QT GWAS for each of the 68 SNPs. Then, the  $PRS_{QT}$  was normalized to a mean of 0 and standard deviation of 1. We used logistic regression to test for association of  $PRS_{QT}$  with case-control status, correcting for PC 1-10. We also used P-value thresholding and R2 pruning with P-values of  $5 \times 10^{-8}$ ,  $1 \times 10^{-5}$ ,  $1 \times 10^{-4}$  and  $1 \times 10^{-3}$ ,  $1 \times 10^{-2}$  and R2 of 0.2 and 0.1 on summary statistics from a European[29] and Japanese[30] descent general population QT-interval GWAS. The resulting 10 models were used to calculate a European and Japanese  $PRS_{QT}$ . The association of  $PRS_{QT}$  with LQTS was assessed using a logistic regression for the European and Japanese cases separately. The best model was selected based on the maximal C-statistic, as recently performed.[31] No other covariate was used to avoid model overfitting.

The odds ratios (ORs) associated with quartile 2, 3 and 4 were calculated using the first  $PRS_{QT}$  quartile as the reference. The association of  $PRS_{QT}$  and known QT predictors with QTc was performed using a univariable linear regression followed by multivariable analysis, including in the final model only those variables with a  $P < 0.05$  in the univariable analyses. The association of  $PRS_{QT}$  quartiles with time to life-threatening arrhythmic events (LAE) was assessed using Cox proportional hazards regression with/without adjustment for classic risk factors. Association analyses of  $PRS_{QT}$  with case-control status, QTc and time to LAE were performed separately in the European and Japanese datasets, followed by a fixed-effects model meta-analysis.

### **Common variant heritability**

We used the generalized restricted maximum likelihood (GREML) approach of GCTA[32] to estimate how much of the variance in LQTS susceptibility could be attributed to common genetic variants (SNP-based heritability,  $h^2_{SNP}$ ). Prior to heritability estimation, we conducted additional stringent genetic QC, as previously suggested (**see Supplementary Methods**).[33] We estimated the SNP-heritability on the liability scale assuming a 0.04% prevalence with

PC1-10 as covariates.[1] We assessed the robustness of heritability estimates from GCTA-GREML using the generalized restricted maximum likelihood (REML) and PCGC (phenotype-correlation genotype-correlation) regression[34] analyses implemented in LDAK[35]. We estimated  $h^2_{\text{SNP}}$  in the overall LQTS and genotype positive LQTS dataset in the both European and Japanese ancestries. Because of small sample size we were not able to estimate  $h^2_{\text{SNP}}$  in genotype negative LQTS patients using the approaches implemented in GCTA or LDAK.

### **Genetic correlation with other traits**

We used bivariate LD score regression[36] to evaluate the genetic correlation between LQTS susceptibility (as obtained in the European descent case-control GWAS) and other cardiac electrical traits[2], namely PR, QRS, QT, heart rate (HR) at rest, HR in response to exercise and recovery, and atrial fibrillation (AF) (see **Supplementary Methods** for origin of summary statistics). We used Bonferroni correction to account for multiple testing ( $P = 0.05 / 7 = 0.0071$ ). We did not constrain the bivariate regression intercepts in any of these analyses given the potential for (modest) sample overlap and population stratification.



## RESULTS

### Clinical characteristics of the case cohort

Demographic and clinical characteristics of the unrelated LQTS cases are presented in **Table IV of the Supplement** separately for the European and Japanese datasets and in **Table 1** for the combined cohort. We included a total of 1,781 unrelated LQTS patients of European (n=1,344, mean QTc  $\pm$  standard deviation (SD): 484 $\pm$ 48ms) and Japanese descent (n=437, QTc: 485 $\pm$ 49ms). A total of 1584 cases (89%) were genotype positive, carrying a rare variant in *KCNQ1* (LQT1, n=800), *KCNH2* (LQT2, n=661), or *SCN5A* (LQT3, n=123), while in 197 (11%) no disease causing variant was identified (i.e. genotype negative) despite extensive genetic testing.

The mean QTc interval in genotype negative cases was higher in comparison to genotype positive ones (500 $\pm$ 52ms vs. 482 $\pm$ 47ms,  $P=2\times 10^{-5}$ ) and in genotype negative cases a family history of SCD <50 years in 1<sup>st</sup> and 2<sup>nd</sup> degree relatives was less frequent compared to genotype positive ones (12.7% vs. 22.9%,  $P=0.001$ ). Of the 1584 genotype positive cases, 1333 (84%) carried an ACMG pathogenic or likely-pathogenic variant and the remainder had a VUS. The QTc did not significantly differ between carriers of VUS and those with a pathogenic or likely pathogenic variant ( $P=0.9$ ).

In total, 429 cases (24%) had a life-threatening arrhythmic event (LAE) at a median age of 28 [17-46] years, with 295 cases (17%) having such an event by age 40. LAE-free survival did not significantly differ between genotype negative and positive cases ( $P=0.8$ ) or between European and Japanese cases ( $P=0.053$ ) (**Figure 1**). In a multivariable Cox proportional hazard model, male sex (OR 1.9;  $P=0.004$ ), QTc>500ms (OR 1.8;  $P=4\times 10^{-6}$ ) and Japanese ancestry (OR 1.4;  $P=0.03$ ) were independent risk factors for LAE (**Table V in the Supplement**). We found a significant sex-puberty interaction ( $P=1\times 10^{-6}$ ), where males were at higher risk of LAE in the pre-pubertal years but lower risk thereafter (**Figure II in the Supplement**). The effect of the conventional risk factors sex ( $P_{\text{interaction}}=0.3$ ) and QTc  $\geq$  500ms ( $P_{\text{interaction}}=0.7$ ) did not differ between genotype positive and genotype negative cases.

Genotype (*KCNQ1*, *KCNH2*, *SCN5A* or negative) significantly affected time to LAE (Logrank test  $P < 0.001$ ; **Figure III in the Supplement**). Cases with a rare variant in *KCNQ1* had a lower risk of LAE compared to *KCNH2*, *SCN5A* and genotype negative ones ( $P < 0.01$  for all comparisons). None of the other post-hoc pairwise comparisons reached statistical significance. Time to LAE did not differ between cases with a VUS and those with pathogenic or likely pathogenic variant (**Figure IV in the Supplement**).

### Case-control GWAS

We conducted a case-control GWAS separately in European (1,238 cases vs. 8,219 controls, genomic test inflation ( $\lambda$ ) = 1.024) and Japanese (418 cases vs. 1,617 controls,  $\lambda$  = 1.034) cases using ancestry-matched controls (**Figure V and VI in the Supplement**), followed by transethnic meta-analysis ( $\lambda$  = 1.028). This uncovered 3 loci reaching the threshold for genome-wide statistical significance (**Table 2, Figure 2, Figure VII and VIII in the Supplement**). The most significant association was obtained for rs12143842 (OR = 1.32, 95% confidence interval (CI): 1.21-1.42,  $P = 1.09 \times 10^{-11}$ ) located upstream of *NOS1AP* (**Figure VIII-a in the Supplement**). The lead SNP at the second locus was located in an intron of *KCNQ1* (rs179405, OR = 1.38, 95%CI: 1.23-1.54,  $P = 1.92 \times 10^{-8}$  (**Figure VIII-b in the Supplement**). At the third locus, the lead SNP, rs17061696 (OR = 1.25, 95%CI: 1.15-1.35,  $P = 4.33 \times 10^{-8}$ ), was located in an intron of *KLF12* (**Figure VIII-c in the Supplement**). All three loci had been previously associated with the QT-interval duration, a measure of myocardial repolarization on the electrocardiogram (ECG), in the general population (**Table 1**).[29] The low-frequency missense variant in *KCNE1*, p.Asp85Asn (rs1805128, OR = 2.78, 95%CI: 1.67-3.90,  $P = 5.31 \times 10^{-7}$ , **Figure VIII-d in the Supplement**) reached the suggestive statistical significance threshold in the European case-control analysis. This variant, which is rare and not well imputed in the Japanese dataset (MAF = 0.001;  $R^2 < 0.3$ ), has the largest reported effect size among the 68 independent SNPs (hereafter referred to as 'QT-SNPs') previously associated with QT-interval in the general population (7.4ms increase per minor allele).[29] Of note, The

KCNE1-p.Asp85Asn variant had a more pronounced effect in genotype negative (OR=7.64, 95%CI: 3.66-15.95,  $P=5.99 \times 10^{-8}$ ) than in genotype positive LQTS (OR=2.28, 95%-CI: 1.46-3.54,  $P=2.59 \times 10^{-4}$ ).

### **Genetic overlap between LQTS and QT-interval in the general population**

The identification of SNPs previously associated with QT-interval in the general population is in line with the fact that QT-interval prolongation on the ECG (representing prolonged cardiac repolarization) is the central intermediate phenotype underlying LQTS. In fact, 23 of the 68 QT-SNPs previously associated with QT-interval in the general population, were associated with LQTS at nominal significance (i.e.  $P < 0.05$ ), while only 4 would be expected under the null hypothesis (**Table VI in the Supplement**). We observed a strong positive correlation between the effect that each of the 68 QT-SNPs had on the QT-interval in the general population[29] and the risk they conferred for LQTS in the current study; this effect was consistent across both the European (**Figure 3a**;  $R^2=0.67$ ;  $P=2.04 \times 10^{-17}$ ) and the Japanese (**Figure 3b**;  $R^2=0.52$ ;  $P=1.43 \times 10^{-10}$ ) datasets. Overlap between genetic risk for LQTS and genetic determinants of the QT-interval in the general population[29] was further demonstrated by genome-wide bivariate LD score regression[36], which detected a significant positive genetic correlation ( $r_g=0.40$ ,  $SE=0.14$ ,  $P=3.2 \times 10^{-4}$ ) between these phenotypes. No significant correlation was found for other cardiac electrical traits (**Figure IX in the Supplement**).

### **Analysis of PRS<sub>QT</sub> in LQTS disease susceptibility**

We then tested the aggregate effect of the 68 QT-SNPs (PRS<sub>QT</sub>) on susceptibility to LQTS by means of polygenic risk score (PRS) analysis (**Table III in the Supplement**). PRS<sub>QT</sub> was significantly associated with a diagnosis of LQTS in the European set, the Japanese set and in the meta-analysis of both datasets (**Figure 4a and 4c**; **Table 3**; meta-analysis  $\beta=0.34$ ,  $SE=0.03$ ,  $P=1.1 \times 10^{-38}$ , heterogeneity  $P=0.15$ ). Similar results were obtained when we excluded common variants located at the known Mendelian LQTS loci from the PRS (**Table VII in the Supplement**). Ten different PRS derived by the pruning and thresholding method on summary

statistics from the European descent general population QT-interval GWAS did not significantly outperform the PRS<sub>QT</sub> in discriminating case-control status (**Table VIII-a in the Supplement**). Similarly, Japanese ancestry-specific PRS derived from summary statistics of a small Japanese QT-interval GWAS[30] had less discriminative accuracy in the Japanese case-control dataset compared to the European-derived PRS<sub>QT</sub>, likely due to the small size of the Japanese QT-interval GWAS (**Table VIII-b in the Supplement**).

We next explored whether the genetic architecture of genotype negative patients (i.e. those lacking a rare variant after extensive genetic testing of the established LQTS disease genes) differed from that of genotype positive patients. This was done by comparing PRS<sub>QT</sub> between both groups, uncovering a significantly higher PRS<sub>QT</sub> in genotype negative patients, pointing to a more prominent role for common variants in disease susceptibility in these patients. This effect was consistently observed in both the European ( $P=5.1 \times 10^{-6}$ , **Figure 4b**) and the Japanese ( $P=2.0 \times 10^{-3}$ , **Figure 4d**) datasets (**Table 3**). Similar results were obtained in a sensitivity analysis correcting for QT-interval, ensuring that enrichment of QT prolonging alleles in the genotype negative patients was not driven by differences in QT-interval ( $P=7.4 \times 10^{-5}$  in Europeans;  $P=2.6 \times 10^{-3}$  in Japanese, **Table 3**). These associations remained statistically significant when we restricted the analysis to patients with a pathogenic or likely pathogenic variant according to American College of Medical Genetics and Genomics and Association of Molecular Pathology (ACMG/AMP) guidelines (i.e. excluding cases with a rare variant of unknown significance (VUS); **Supplementary Methods, Table IX and X in the Supplement**). Increasing PRS<sub>QT</sub> quartiles were associated with a significantly higher disease susceptibility for genotype negative LQTS compared to the lowest quartile (**Figure 5, Table XI in the Supplement**). Notably, using a PRS<sub>QT</sub> percentile threshold of 80, 90 and 95, individuals above the threshold compared to those below have an OR [95%CI] of 2.9 [2.2-4.0], 4.1 [2.9-5.8] and 5.7 [3.9-8.4], respectively, for genotype negative LQTS. Of interest, the higher PRS<sub>QT</sub> in genotype negative patients compared to genotype positive patients was reflected by the larger difference in PRS<sub>QT</sub> between genotype negative patients vs. controls (**Table 3**; meta-

analysis  $\beta=0.735$ ,  $SE=0.074$ ,  $P=2.24 \times 10^{-23}$ ) compared to genotype positive vs. controls (**Table 3**; meta-analysis  $\beta=0.294$ ,  $SE=0.028$ ,  $P=1.09 \times 10^{-25}$ ).

### **Common variant heritability of LQTS**

To evaluate the proportion of variance in LQTS susceptibility explained by common genetic variants ( $h^2_{\text{SNP}}$ ) we used the generalized restricted maximum likelihood (GREML) approach of GCTA.[32,37] Assuming a disease prevalence of 0.04%, [1] the SNP heritability estimate on the liability scale was  $h^2_{\text{SNP}} = 0.148$  ( $SE=0.019$ , 95%-CI: 0.111-0.185,  $P=5.0 \times 10^{-18}$ ) in the overall European LQTS dataset.  $h^2_{\text{SNP}}$  was similar when the analysis was restricted to genotype positive LQTS patients. Similar results were also observed in the Japanese dataset and when using the PCGC (phenotype-correlation genotype-correlation) regression[34] and the GREML estimation implemented in LDAK[35], as well as when we restricted  $h^2_{\text{SNP}}$  analyses to only patients with a pathogenic or likely pathogenic variant (**Table XII in the Supplement**).

### **Association of single SNPs and PRS<sub>QT</sub> with LQTS severity**

To identify genetic modifiers of disease severity we conducted a GWAS for QT-interval within the LQTS cases which did not uncover any genome-wide significant loci (**Figure X in the Supplement**). None of the 68 SNPs previously associated with QTc in the general population[29] showed association with QTc after Bonferroni correction. PRS<sub>QT</sub> showed a weak positive correlation with QTc in the European cases (correlation coefficient [ $r$ ]=0.06;  $P=0.042$ ; **Figure XI in the Supplement**). In a multivariable linear regression model including clinical covariates associated with QTc (age at ECG recording, LQTS type and sex), PRS<sub>QT</sub> was not significantly associated with QTc (**Table XIII in the Supplement**). Similarly, in a sub-analysis restricted to probands (comprising 97% of the total of unrelated LQTS cases) using the multivariable linear regression model, PRS<sub>QT</sub> was not significantly associated with QTc (**data not shown**). In exploratory subgroup analyses, PRS<sub>QT</sub> was independently associated with QTc in *KCNH2* rare variant carriers but not in *KCNQ1* rare variant carriers (**Table XIII in**

**the Supplement**). This result was not replicated in the Japanese LQTS dataset.  $PRS_{QT}$  was not significantly associated with time to LAE in neither Europeans nor Japanese cases (**Figure XII in the Supplement**).

## DISCUSSION

Our findings establish an important role for common genetic variation in LQTS susceptibility and support a complex (polygenic) architecture in genotype negative LQTS. Case-control GWAS identified 3 genome-wide significant risk loci near *NOS1AP*, *KCNQ1* and *KLF12*. Heritability analysis demonstrated that ~15% of LQTS disease liability is attributable to common genetic variation. Polygenic risk score analysis testing the aggregate effect of SNPs previously associated with QT interval in the general population ( $PRS_{QT}$ ) identified a higher  $PRS_{QT}$  in LQTS cases compared to controls and higher  $PRS_{QT}$  in genotype negative vs. genotype positive LQTS.

### Shared genetics of LQTS and QT interval in the general population

The case-control GWAS uncovered three genetic LQTS susceptibility loci at genome-wide statistical significance near *NOS1AP*, *KCNQ1* and *KLF12*, and one missense variant in *KCNE1* at the suggestive threshold (**Figure 2**). The association of SNPs at *KCNQ1* points to the involvement of common variants acting alongside rare variants in these genes in mediating disease susceptibility, akin to what was previously reported for common and rare variation in and around the *SCN5A* gene in Brugada syndrome.[16] All 4 risk loci had been previously implicated in genetic control of the QT interval by GWAS in the general population.[29] For the 68 SNPs associated with QT interval in the general population, we noted a strong positive correlation between their effect on QT interval (obtained in the general population) and their odds ratio for LQTS susceptibility, indicating, as expected, that the larger the effect a SNP has on the QT interval, the more it increases LQTS susceptibility (**Figure 3**). The strong genetic correlation between LQTS susceptibility and QT interval in the general population provides quantitative support for genetic overlap (**Figure IX in the Supplement**).

The association with the highest effect in the case-control GWAS was found for the p.Asp85Asn missense variant in *KCNE1* (rs1805128). This variant increased susceptibility for LQTS in the overall cohort, but had a more prominent effect in genotype negative LQTS with

an odds ratio of ~7 (versus an odds ratio of 2 in genotype positive patients). This variant has an allele frequency of ~1.2% in non-Finnish Europeans, and ~0.5% in East Asians and has the largest effect size on the QT in the general (European descent) population (7.42 ms per minor [Asn] allele).[29] It has been shown to be enriched in patients with drug-induced torsades de pointes[38].

### **Genetic architecture of genotype positive LQTS**

LQTS has traditionally been seen as a monogenic disorder mostly attributed to a rare variant with a drastic effect on ion channel function. We now demonstrate that a considerable extent (~15%) of disease liability is attributable to common genetic variation. In genotype positive LQTS families, where the penetrance of pathogenic variants may be low for certain variants[8], the contribution of common variants to disease susceptibility may also contribute to variable disease penetrance. It has been well established that LQTS probands have a longer QT interval and greater arrhythmic risk compared to family members carrying the same variant.[7,13,39] This observed increased penetrance in probands may result from a greater burden of common QT-prolonging variants compared to other, less-severely affected or unaffected mutation-carriers. However, since this study was comprised of only unrelated patients, this remains to be determined. Whether the  $PRS_{QT}$  could discriminate between affected vs. unaffected mutation carrier family members is intuitive appealing but remains to be formally demonstrated.

### **Genotype negative LQTS: A polygenic subtype of LQTS?**

Polygenic risk score analysis testing the aggregate effect of SNPs previously associated with QT interval in the general population ( $PRS_{QT}$ ) identified a higher  $PRS_{QT}$  in genotype negative vs. genotype positive patients. This observation points to genotype negative LQTS, comprising ~10% of LQTS patients, as a polygenic subtype of the disorder where the underlying etiology involves, at least in part, a high burden of common QT prolonging alleles. As such, genetic susceptibility in genotype negative patients may not be determined to a large extent by one



strong genetic factor as occurs in genotype positive patients, but results from the accumulation of multiple variants (polygenic inheritance). The lower rate of family history of SCD in genotype negative LQTS patients is in line with polygenic inheritance. Our observations corroborate findings in other heritable phenotypes, such as familial hypercholesterolemia (FH), where patients without a disease-causing variant in the *LDLR*, *APOB* and *PCSK9* genes have a higher PRS based on LDL modulating variants in comparison to those with rare FH causing genetic variants.[40] As such, the accumulation of multiple discrete common variants may confer risk similar to a monogenic mutation. This was recently demonstrated for common disorders such as coronary artery disease and atrial fibrillation, where individuals at the upper extreme of the PRS distribution had a risk of developing the disease reportedly comparable to carriers of a monogenic mutation.[31] The overlap in the PRS<sub>QT</sub> distributions among genotype negative LQTS cases and controls (**Figure 4**) suggests that other factors are involved, possibly including low-frequency genetic variants with intermediate effect sizes as well as other common variants with smaller effect sizes.

In addition to providing insight into the genetic architecture of genotype negative LQTS, we here also describe for the first time the natural history in these patients. All ~200 genotype negative LQTS patients met diagnostic criteria for definite LQTS (i.e. QTc>500ms or LQTS score  $\geq 3.5$ ) and underwent sequencing of the unequivocal non-syndromic LQTS genes. Patients with genotype negative LQTS had a higher QTc in comparison to LQT1-3 patients but similar event-free survival as their genotype positive counterparts (**Figure 1**). The effect of established clinical risk factors, i.e. sex and QTc-duration, did not significantly differ between genotype positive and negative (no interaction effect) suggesting they may also be used to stratify risk of events in genotype negative LQTS.

### **Common variants do not contribute to LQTS severity within probands**

We sought to identify genetic modifiers of LQTS. In contrast to the case-control GWAS, GWAS for QTc and arrhythmic events within the unrelated LQTS cases did not uncover any genome-wide significant locus. PRS<sub>QT</sub> was also not significantly associated with QTc nor with the occurrence of events. At first glance, this may seem contradictory to previous studies in LQTS that demonstrated a modulatory effect of SNPs at *NOS1AP* on the QTc and arrhythmic events[10,11,13], as well as a study in the general population that showed a modulatory effect of PRS derived from prior GWAS on QT-interval.[41] For example, a study we previously conducted in LQT2 patients uncovered strong associations with large effect sizes (>12 ms/allele) for SNPs at *NOS1AP*. [13] An important difference however, is that the current study did not include family members but only one patient per family (97% probands), whereas the previous studies considered both probands and genotype positive relatives. Conceptually, inclusion of both probands and relatives results in greater variation in QTc and is thus expected to increase statistical power for detection of modulatory effects. Moreover, the different rare variants in the patients we studied here are associated with biophysical defects of varying severity. As such, they are also expected to contribute to interindividual variability which is difficult to account for. For instance, LQT2 patients with pore-region variants are known to be more severely affected than other LQT2 patients.[42] Indeed, in a sub-analysis, restricted to European LQT2 patients, where we accounted for the mutation location (pore versus non-pore), we detected an association of PRS<sub>QT</sub> with QTc. In sum, our data show that common variants do not affect disease severity across all probands studied. Further studies are needed to explore their predictive role in family members.

### **Potential clinical implications**

In genotype negative LQTS, disease susceptibility estimation for relatives does not follow a Mendelian pattern. In our cohort, a positive family history of SCD was less often observed in genotype negative individuals compared to genotype positive ones, suggesting that risk for family members in genotype negative patients may be lower. Polygenicity in genotype negative

individuals implies that risk is not primarily attributable to one genetic factor inherited from one of the biological parents as is the case for autosomal dominant LQTS. In such cases, cascade screening may necessitate clinical evaluation of both maternal and paternal family members. Future clinical utility of genetic testing based on polygenic inheritance necessitates the availability of polygenic risk scores with high discriminative capacity. The discriminative capacity of a PRS based on QT modulating SNPs is expected to improve as knowledge concerning variants that modulate the QT interval become known, for example through larger GWAS studies, or by combining it with non-genetic modifiers. In a recent study, a PRS based on 61 QT SNPs (a subset of the 68 QT modulating SNPs included in the PRS<sub>QT</sub> used herein) explained a substantial proportion of QT interval response to QT prolonging drugs in a trial of 3 QT-prolonging drugs conducted in healthy individuals, as well as risk of torsade de pointes in a case-control study.[43] This provides further support to a liability threshold model whereby multiple factors, genetic and non-genetic, impact on cardiac repolarization and determine arrhythmic risk. In this respect, calculation of PRS<sub>QT</sub> for the purpose of preventive avoidance of QT-prolonging drugs may be desirable for relatives of genotype negative LQTS. Clearly, further studies are needed to address how testing for polygenic susceptibility may become clinically useful.

### **Study limitations**

Although in genotype negative LQTS patients we performed sequencing of the coding region of non-syndromic LQTS genes, this may have missed copy number variation or disease-causing variants in the non-coding region[44] of established genes as well as mutations in yet unknown disease genes. This may have blunted the differences between genotype negative and genotype patients and thus would not affect the study conclusions. Despite being the largest international dataset of unrelated LQTS patients published to date, the study had limited statistical power to detect lower effect associations at GWAS significance threshold. The prespecified design of meta-analysing European and Japanese GWAS may also miss disease loci with differences in haplotype structure among European and East Asian chromosomes.

Nonetheless, GWAS in separate ancestries did not detect any association at GWAS threshold. Finally, studies in larger patient sets are required to further refine our understanding of the genetic architecture underlying LQTS in genotype negative patients.

## **Conclusion**

This work establishes an important role for common genetic variation in susceptibility to LQTS. Common genetic variation affecting the QT interval in the general population contributes to disease susceptibility in both genotype positive and genotype negative LQTS. The role of common variants is predominant in genotype negative LQTS, suggesting that the latter may constitute a polygenic form of LQTS. Increasing burden of QT-prolonging common variants (e.g.  $PRS_{QT}$ ) is associated with higher susceptibility for LQTS but is not associated with disease severity within LQTS probands. Further studies are needed to assess the role of polygenic risk within LQTS families.

## **ACKNOWLEDGEMENTS**

The authors thank Estelle Baron, Simon Lecointe, Annabelle Rajalu, Aurélie Thollet and Florence Kyndt for their help in DNA sample preparation and regulatory processes at the Nantes centre, the Nantes biological resource centre for biobanking (CHU Nantes, Hôtel Dieu, Centre de ressources biologiques (CRB), Nantes, F-44093, France (BRIF: BB-0033-00040)), and the National Referral Centre for Inherited Cardiac Arrhythmias of Nantes and its associated competence centres. We thank Prof. Pier D. Lambiase and Prof. Patricia B. Munroe for sharing summary statistics. We thank Dr. Nicky Whiffin for help with using CardioClassifier. The KORA research platform (KORA, Cooperative Research in the Region of Augsburg) was initiated and financed by the Helmholtz Zentrum München - German Research Center for Environmental Health, which is funded by the German Federal Ministry of Education and Research and by the State of Bavaria. Furthermore, KORA research was supported within the Munich Center of Health Sciences (MC Health), Ludwig-Maximilians-Universität, as part of LMUinnovativ. This study makes use of data generated by the Wellcome Trust Case-Control Consortium. A full list of the investigators who contributed to the generation of the data is available from [www.wtccc.org.uk](http://www.wtccc.org.uk). Funding for the project was provided by the Wellcome Trust under award 076113, 085475 and 090355.

## **SOURCES OF FUNDING**

CRB, NL, AAW, JPVt and HLT acknowledge the support from the Dutch Heart Foundation (CVON 2018-30 PREDICT2 project to CRB, HLT, and AAW) and the Netherlands Organization for Scientific Research (VICI fellowship, 016.150.610, to CRB). HLT has received funding from the European Union's Horizon 2020 research and innovation programme under acronym ESCAPE-NET (The European Sudden Cardiac Arrest network towards Prevention, Education, New Effective Treatment), registered under grant agreement No 733381. RT received support from the Canadian Heart Rhythm Society (George Mines Award), the European Society of Cardiology (research award), and the Philippa and Marvin Carsley Cardiology Chair and is currently a clinical research scholar of the Fonds de Recherche du Québec—Santé; PJS and

LC acknowledge the support of the Leducq Foundation for cardiovascular research grant 18CVD05: Towards Precision Medicine with Human iPSCs for Cardiac Channelopathies. VP and JBG were supported by a grant from Hopitaux Universitaires du Grand Ouest and Fondation Maladies Rares [RC17\_0357]; JB was supported by the H2020-MSCA-IF-2014 Program of the European Commission (RISTRAD-661617) and by the Regional Council of Pays-de-la-Loire (Etoile montante: REGIOCARD); JJS was supported by the Fondation pour la Recherche Médicale (DEQ20140329545) and by the National Agency for Research (ANR-GENSUD-14-CE10-0001); AR and UBD are supported by The Swedish Heart-Lung Foundation; ERB is supported by the Higher Education Funding Council for England and the British Heart Foundation (BHF) and acknowledges support from Cardiac Risk in the Young. YDW had received support through an Academic Clinical Fellowship from the National Institute of Health Research. YDW and ERB gratefully acknowledge funding and ongoing support from the James Lancaster Memorial Fund sponsored by McColl's RG Ltd.; This work was partly supported by a Health and Labour Sciences Research Grant from the Ministry of Health, Labour and Welfare of Japan (H22-032, H24-033, H26-040, and H29-055) to MY; NM is supported by: AMED (19kk0305011h0001); WS and TA acknowledge support from a Health Science Research Grant from the Ministry of Health, Labor and Welfare of Japan for Clinical Research on Measures for Intractable Diseases (H24-033, H26-040, and H27-032). EA is supported by Instituto de Salud Carlos III (FIS PI16/01203 and PI17/01690) co-funded by ERDF/ESF, "Investing in Your Future"; This work was supported by the John and Birthe Meyer Foundation, The Hallas-Møller Emerging Investigator Novo Nordisk (NNF17OC0031204); The iPSYCH study was funded by the Lundbeck Foundation Initiative for Integrative Psychiatric Research. This research has been conducted using the Danish National Biobank resource; The Cardiac Inherited Disease Registry has been supported by Cure Kids; AW is supported by the Hugh Green Foundation; PTE is supported by the Fondation Leducq (14CVD01), by grants from the National Institutes of Health (1R01HL092577, R01HL128914, K24HL105780); and the American Heart Association (18SFRN34110082).; SAL is supported by NIH grant 1R01HL139731 and American Heart Association 18SFRN34250007; BL is supported by an

European Research Council (ERC) consolidator grant; DMR is supported by P50 GM115305 and MBS by K23 HL127704; CA is supported by NIH - HL47678 and HL138103. PGP is supported by The Swedish Heart-Lung Foundation, grant #20180444. KLT, DJT, JMB, and MJA were supported by the Mayo Clinic Windland Smith Rice Comprehensive Sudden Cardiac Death Program.

## **DISCLOSURES**

ERB received prior research funds from Biotronik and consulting for Medtronic; EA received speaking fees from Biosense Webster; SAL receives sponsored research support from Bristol Myers Squibb / Pfizer, Bayer AG, and Boehringer Ingelheim, and has consulted for Bristol Myers Squibb / Pfizer and Bayer AG; PTE is supported by a grant from Bayer AG to the Broad Institute focused on the genetics and therapeutics of cardiovascular diseases. PTE has also served on advisory boards or consulted for Bayer AG, Quest Diagnostics, and Novartis. DC is a scientific consultant for Bio4Dreams. MJA is a consultant for Audentes Therapeutics, Boston Scientific, Gilead Sciences, Invitae, Medtronic, Myokardia, and St. Jude Medical. MJA and Mayo Clinic have an equity/royalty-based licensing agreement with AliveCor.

## **Supplemental Materials**


Expanded Methods

Online-only Figures I-XII

Supplemental Excel File with Supplementary Table I-XIII

References [45–57] (45-57)


**Figure 1** | Kaplan-Meier LAE-free survival curves stratified by ancestry






**Figure 1 | Kaplan-Meier LAE-free survival curves stratified by ancestry.** EU and JP refer to European and Japanese LQTS cases, respectively. Geno+ and Geno- refer to genotype positive and genotype negative LQTS cases, respectively. LAE, life-threatening arrhythmic event defined as the composite of out of hospital cardiac arrest or hemodynamically unstable ventricular tachycardia/fibrillation. Logrank test  $P=0.3$ .

Figure 2 | Manhattan plot of LQTS case-control meta-analysis




**Figure 2 | Manhattan plot of LQTS case-control meta-analysis.** Manhattan plot displaying the base-pair position of each of the tested SNPs (each dot represents an individual SNP) along the chromosomes on the x axis and the corresponding  $-\log_{10}$  transformed association  $P$  value on the y axis. The association  $P$ -values from the meta-analysis of the two GWAS conducted separately in European and Japanese cases and controls, respectively, are displayed. The upper and lower dashed lines indicate the genome-wide significance ( $P < 5 \times 10^{-8}$ ) and suggestive significance ( $P < 1 \times 10^{-6}$ ) thresholds, respectively. SNPs at genomic regions that reached the genome-wide or suggestive significance thresholds, are marked in red, whereas SNPs from other regions are marked in black or grey. The association for variant rs1805128 (*KCNE1*:p.Asp85Asn) is solely driven by the European analysis since it is not well imputed and rare ( $R^2 < 0.3$ . MAF=0.001) in the Japanese dataset.

**Figure 3 | Correlation of effect size of QT-associated SNPs with their effect size in LQTS GWAS**




**Figure 3 | Correlation of effect size of QT-associated SNPs with their effect size in LQTS GWAS.** The x-axis represents the effect estimates from the QT-interval GWAS conducted in the general population (milliseconds per alternative allele) and the y-axis the effect of each of these QT-interval associated alleles on disease risk of LQTS [ $\ln(\text{OR})$ ] in the European (**A, blue**) and Japanese (**B, red**) datasets. All 68 SNPs associated with QT in the general population were assessed in Europeans, whereas 60 SNPs were properly imputed in the Japanese dataset. In the LQTS-GWAS meta-analysis, 23/68 SNPs previously associated with the QT in the general population reached nominal significance (see **Supplementary Table 6**). Loci that reached genome-wide significance in the LQTS cases-control transethnic meta-analysis (*NOS1AP*-rs12143842, *KCNQ1*-rs179405 and *KLF12*-rs728926) and *KCNE1*-rs1805128 are identified with text.

**Figure 4 |** Distribution of PRS<sub>QT</sub> in controls, all LQTS and genotype positive and negative subgroups



**Figure 4 | Distribution of PRS<sub>QT</sub> in controls, LQTS and genotype positive and negative subgroups.** The X-axis represents the QT polygenic score (PRS<sub>QT</sub>) in the European (**A-B**, blue) and Japanese (**C-D**, red) LQTS case-control datasets. In **A and C**, all LQTS cases are grouped regardless of whether they are genotype positive or negative, whereas in **B and D** cases have been stratified in genotype positive and negative LQTS subgroups. PRS<sub>QT</sub> was normalized to a mean of 0 and standard deviation of 1. Reported P values refer to the effect of PRS<sub>QT</sub> in a logistic regression correcting for the first 10 principal components. \*Refers to case-control association. Comparison of PRS<sub>QT</sub> between genotype negative vs. genotype positive LQTS uncovered a significantly higher PRS<sub>QT</sub> in genotype negative patients. This effect was consistently observed in both the European ( $P=5.1 \times 10^{-6}$ , **Figure 4b**) and the Japanese ( $P=2.0 \times 10^{-3}$ , **Figure 4d**) patients (**Table 3**).

**Figure 5** | Increasing genotype negative LQTS risk with increasing PRS<sub>QT</sub> quartiles





**Figure 5 | Increasing LQTS risk with increasing PRS<sub>QT</sub> quartiles.** Odds ratio (OR) for genotype negative LQTS (filled circles) and 95% confidence intervals (vertical bars) associated with each PRS<sub>QT</sub> quartile taking the first PRS<sub>QT</sub> quartile as the reference. Data shown correspond to a meta-analysis of effects computed separately in the European and Japanese datasets. P-values refer to comparison of each quartile against the first quartile.

**Table 1** | Clinical characteristics of all unrelated LQTS cases

Parameter	Genotype positive (n=1584)	Genotype negative (n=197)
Male, no. (%)	584/1584 (37)	76/197 (39)
QTc mean $\pm$ SD, ms	482 $\pm$ 47	500 $\pm$ 52
Genotype, no. (%)		
<i>KCNQ1</i>	800/1584 (50)	-
<i>KCNH2</i>	661/1584 (42)	-
<i>SCN5A</i>	123/1584 (7.8)	-
Syncope, no. (%)	722/1584 (46)	75/197 (38)
LAE (OHCA or VF/VT) before age 41, no. (%)	262/1578 (17)	33/196 (17)
Age at first LAE, median [IQR]	21 [13-29]	26 [16-35]
Treatment during follow-up, no. (%)		
Beta-blocker	1169/1487 (79)	124/168 (74)
ICD	277/1562 (18)	38/172 (22)
PM	50/1565 (3.2)	11/171 (6.4)
LCSD	29/1583 (1.8)	1/171 (0.6)
Family history of SCD <50 years of age, no. (%)	323/1409 (23)	24/189 (13)

**Table legend:**

ICD, implantable cardioverter-defibrillator; IQR, interquartile range; LAE, life-threatening arrhythmic event; LCSD, left cardiac sympathetic denervation; OHCA, out of hospital cardiac arrest; PM, pacemaker; SCD, sudden cardiac death, SD, standard deviation; VF/VT, hemodynamically unstable ventricular fibrillation/tachycardia

**Table 2 |** Significant loci in LQTS case-control GWAS

Lead SNP	GRCh37	Alternative allele	Reference allele	Closest gene	Meta-analysis			European			Japanese			Effect on QT (ms) *
					OR	95%CI	P	AAF (controls /cases)	OR	P	AAF (controls /cases)	OR	P	
rs12143842	1:162033890	T	C	<i>NOS1AP</i>	1.31	1.21-1.42	1.09E-11	0.26/0.32	1.29	7.34E-08	0.38/0.47	1.41	2.13E-05	3.5
rs179405	11:2525395	A	G	<i>KCNQ1</i>	1.38	1.23-1.54	1.92E-08	0.14/0.17	1.34	4.03E-06	0.10/0.14	1.63	5.42E-04	1.9 <sup>#</sup>
rs17061696	13:74511991	C	G	<i>KLF12</i>	1.25	1.15-1.35	4.33E-08	0.37/0.43	1.27	8.91E-08	0.19/0.21	1.16	1.43E-01	0.58

**Table legend:** 95%CI, 95% confidence interval; AAF, alternative allele frequency; GRCh37, genomic position on build GRCh37; OR, odds ratio per alternative allele. \* QT increase (in ms) per alternative allele in the general population, # The lead SNP at the *KCNQ1* locus (rs179405) is in linkage disequilibrium (LD) with rs7122937 ( $R^2=0.497$ ) which had been previously associated with QT-interval in the general population (1.9 ms increase per risk allele).

**Table 3 |** Association of PRS<sub>QT</sub> with LQTS

	European				Japanese				Meta-analysis			
Association analysis of PRS <sub>QT</sub>	n	$\beta$	SE	P	n	$\beta$	SE	P	n	$\beta$	SE	P
All LQTS vs. Controls	1,238/ 8,219	0.322	0.030	4.93E-26	418/ 1,671	0.412	0.055	6.16E-14	1,656/ 9,890	0.343	0.0263	1.08E-38
Genotype positive LQTS vs. Controls	1,115/ 8,219	0.277	0.032	3.47E-18	356/ 1,671	0.348	0.058	2.52E-09	1,471/ 9,890	0.294	0.028	1.09E-25
Genotype negative LQTS vs. Controls	123/ 8,219	0.733	0.090	3.74E-16	62/ 1,671	0.740	0.129	1.19E-08	185/ 9,890	0.735	0.0738	2.24E-23
Genotype negative vs. Genotype positive LQTS	123/ 1,115	0.447	0.098	5.05E-06	62/ 356	0.401	0.129	2.01E-03	185/ 1,471	0.430	0.078	3.54E-08
Genotype negative vs. Genotype positive LQTS*	123/ 1,115	0.409	0.103	7.36E-05	62/ 356	0.393	0.130	2.62E-03	185/ 1,471	0.403	0.0807	6.05E-07

**Table legend:**  $\beta$ , regression coefficient; n, sample size (cases/controls); P, P value; SE, standard error; \*Correcting for QTc

## URLs

LDAK: <http://dougsspeed.com/ldak>

LD score regression: <https://github.com/bulik/ldsc>

FUMA: <http://fuma.ctglab.nl>

PLINK: <https://www.cog-genomics.org/plink2>

METAL: <https://genome.sph.umich.edu/wiki/METAL>

Michigan Imputation Server: <https://imputationserver.sph.umich.edu>

## REFERENCES

- [1] Schwartz PJ, Stramba-Badiale M, Crotti L, Pedrazzini M, Besana A, Bosi G, et al. Prevalence of the congenital long-QT syndrome. *Circulation* 2009;120:1761–1767. doi:10.1161/CIRCULATIONAHA.109.863209.
- [2] Bezzina CR, Lahrouchi N, Priori SG. Genetics of Sudden Cardiac Death. *Circ Res* 2015;116:1919–1936. doi:10.1161/CIRCRESAHA.116.304030.
- [3] Tester DJ, Ackerman MJ. Postmortem long QT syndrome genetic testing for sudden unexplained death in the young. *J Am Coll Cardiol* 2007;49:240–246. doi:10.1016/j.jacc.2006.10.010.
- [4] Lahrouchi N, Raju H, Lodder EM, Papatheodorou E, Ware JS, Papadakis M, et al. Utility of Post-Mortem Genetic Testing in Cases of Sudden Arrhythmic Death Syndrome. *J Am Coll Cardiol* 2017;69:2134–2145. doi:10.1016/j.jacc.2017.02.046.
- [5] Priori SG, Wilde AA, Horie M, Cho Y, Behr ER, Berul C, et al. HRS/EHRA/APHRS expert consensus statement on the diagnosis and management of patients with inherited primary arrhythmia syndromes: document endorsed by HRS, EHRA, and APHRS in May 2013 and by ACCF, AHA, PACES, and AEPC in June 2013. *Heart Rhythm* 2013;10:1932–1963. doi:10.1016/j.hrthm.2013.05.014.
- [6] Priori SG, Napolitano C, Schwartz PJ. Low penetrance in the long-QT syndrome: clinical impact. *Circulation* 1999;99:529–533.
- [7] Goldenberg I, Horr S, Moss AJ, Lopes CM, Barsheshet A, McNitt S, et al. Risk for life-threatening cardiac events in patients with genotype-confirmed long-QT syndrome and normal-range corrected QT intervals. *J Am Coll Cardiol* 2011;57:51–59. doi:10.1016/j.jacc.2010.07.038.
- [8] Mazzanti A, Maragna R, Vacanti G, Monteforte N, Bloise R, Marino M, et al. Interplay Between Genetic Substrate, QTc Duration, and Arrhythmia Risk in Patients With Long

- QT Syndrome. *J Am Coll Cardiol* 2018;71:1663–1671. doi:10.1016/j.jacc.2018.01.078.
- [9] Priori SG, Schwartz PJ, Napolitano C, Bloise R, Ronchetti E, Grillo M, et al. Risk stratification in the long-QT syndrome. *N Engl J Med* 2003;348:1866–1874. doi:10.1056/NEJMoa022147.
- [10] Crotti L, Monti MC, Insolia R, Peljto A, Goosen A, Brink PA, et al. *NOS1AP* Is a Genetic Modifier of the Long-QT Syndrome. *Circulation* 2009;120:1657–1663. doi:10.1161/CIRCULATIONAHA.109.879643.
- [11] Tomás M, Napolitano C, De Giuli L, Bloise R, Subirana I, Malovini A, et al. Polymorphisms in the *NOS1AP* gene modulate QT interval duration and risk of arrhythmias in the long QT syndrome. *J Am Coll Cardiol* 2010;55:2745–2752. doi:10.1016/j.jacc.2009.12.065.
- [12] Duchatelet S, Crotti L, Peat RA, Denjoy I, Itoh H, Berthet M, et al. Identification of a *KCNQ1* Polymorphism Acting as a Protective Modifier Against Arrhythmic Risk in Long-QT Syndrome. *Circ Cardiovasc Genet* 2013;6:354–361. doi:10.1161/CIRCGENETICS.113.000023.
- [13] Kolder ICRM, Tanck MWT, Postema PG, Barc J, Sinner MF, Zumhagen S, et al. Analysis for Genetic Modifiers of Disease Severity in Patients with Long QT Syndrome Type 2. *Circ Cardiovasc Genet* 2015. doi:10.1161/CIRCGENETICS.114.000785.
- [14] Schwartz PJ, Crotti L, George AL. Modifier genes for sudden cardiac death. *Eur Heart J* 2018;39:3925–3931. doi:10.1093/eurheartj/ehy502.
- [15] Ackerman MJ, Priori SG, Willems S, Berul C, Brugada R, Calkins H, et al. HRS/EHRA Expert Consensus Statement on the State of Genetic Testing for the Channelopathies and Cardiomyopathies: This document was developed as a partnership between the Heart Rhythm Society (HRS) and the European Heart Rhythm Association (EHRA). *Europace* 2011;13:1077–1109. doi:10.1093/europace/eur245.

- [16] Bezzina CR, Barc J, Mizusawa Y, Remme CA, Gourraud J-B, Simonet F, et al. Common variants at SCN5A-SCN10A and HEY2 are associated with Brugada syndrome, a rare disease with high risk of sudden cardiac death. *Nat Genet* 2013;45:1044–1049. doi:10.1038/ng.2712.
- [17] Adler A, Novelli V, Amin AS, Abiusi E, Care M, Nannenberg EA, et al. An International, Multicentered, Evidence-Based Reappraisal of Genes Reported to Cause Congenital Long QT Syndrome. *Circulation* 2020;141:418–428. doi:10.1161/CIRCULATIONAHA.119.043132.
- [18] Giudicessi JR, Rohatgi RK, Tester DJ, Ackerman MJ. Variant Frequency and Clinical Phenotype Call Into Question the Nature of Minor, Nonsyndromic Long-QT Syndrome-Susceptibility Gene-Disease Associations. *Circulation* 2020;141:495–497. doi:10.1161/CIRCULATIONAHA.119.043131.
- [19] Roberts JD, Asaki SY, Mazzanti A, Bos JM, Tuleta I, Muir AR, et al. An International Multicenter Evaluation of Type 5 Long QT Syndrome: A Low Penetrant Primary Arrhythmic Condition. *Circulation* 2020;141:429–439. doi:10.1161/CIRCULATIONAHA.119.043114.
- [20] Lek M, Karczewski KJ, Minikel E V., Samocha KE, Banks E, Fennell T, et al. Analysis of protein-coding genetic variation in 60,706 humans. *Nature* 2016;536:285–291. doi:10.1038/nature19057.
- [21] Kobayashi Y, Yang S, Nykamp K, Garcia J, Lincoln SE, Topper SE. Pathogenic variant burden in the ExAC database: an empirical approach to evaluating population data for clinical variant interpretation. *Genome Med* 2017;9:13. doi:10.1186/s13073-017-0403-7.
- [22] Walsh R, Thomson KL, Ware JS, Funke BH, Woodley J, McGuire KJ, et al. Reassessment of Mendelian gene pathogenicity using 7,855 cardiomyopathy cases



and 60,706 reference samples. *Genet Med* 2016. doi:10.1038/gim.2016.90.

- [23] Richards S, Aziz N, Bale S, Bick D, Das S, Gastier-Foster J, et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med* 2015;17:405–423. doi:10.1038/gim.2015.30.
- [24] Vink AS, Neumann B, Lieve KVV, Sinner MF, Hofman N, el Kadi S, et al. Determination and Interpretation of the QT Interval. *Circulation* 2018;138:2345–2358. doi:10.1161/CIRCULATIONAHA.118.033943.
- [25] Anderson CA, Pettersson FH, Clarke GM, Cardon LR, Morris AP, Zondervan KT. Data quality control in genetic case-control association studies. *Nat Protoc* 2010;5:1564–1573. doi:10.1038/nprot.2010.116.
- [26] Das S, Forer L, Schönherr S, Sidore C, Locke AE, Kwong A, et al. Next-generation genotype imputation service and methods. *Nat Genet* 2016;48:1284–1287. doi:10.1038/ng.3656.
- [27] Willer CJ, Li Y, Abecasis GR. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics* 2010;26:2190–2191. doi:10.1093/bioinformatics/btq340.
- [28] Watanabe K, Taskesen E, van Bochoven A, Posthuma D. Functional mapping and annotation of genetic associations with FUMA. *Nat Commun* 2017;8:1826. doi:10.1038/s41467-017-01261-5.
- [29] Arking DE, Pulit SL, Crotti L, van der Harst P, Munroe PB, Koopmann TT, et al. Genetic association study of QT interval highlights role for calcium signaling pathways in myocardial repolarization. *Nat Genet* 2014;46:826–836. doi:10.1038/ng.3014.
- [30] Sano M, Kamitsuji S, Kamatani N, Hong K-W, Han B-G, Kim Y, et al. Genome-wide

- association study of electrocardiographic parameters identifies a new association for PR interval and confirms previously reported associations. *Hum Mol Genet* 2014;23:6668–6676. doi:10.1093/hmg/ddu375.
- [31] Khera A V., Chaffin M, Aragam KG, Haas ME, Roselli C, Choi SH, et al. Genome-wide polygenic scores for common diseases identify individuals with risk equivalent to monogenic mutations. *Nat Genet* 2018;50:1219–1224. doi:10.1038/s41588-018-0183-z.
  - [32] Yang J, Lee SH, Goddard ME, Visscher PM. GCTA: a tool for genome-wide complex trait analysis. *Am J Hum Genet* 2011;88:76–82. doi:10.1016/j.ajhg.2010.11.011.
  - [33] Yang J, Zeng J, Goddard ME, Wray NR, Visscher PM. Concepts, estimation and interpretation of SNP-based heritability. *Nat Genet* 2017;49:1304–1310. doi:10.1038/ng.3941.
  - [34] Golan D, Lander ES, Rosset S. Measuring missing heritability: inferring the contribution of common variants. *Proc Natl Acad Sci U S A* 2014;111:E5272-81. doi:10.1073/pnas.1419064111.
  - [35] Weissbrod O, Flint J, Rosset S. Estimating SNP-Based Heritability and Genetic Correlation in Case-Control Studies Directly and with Summary Statistics. *Am J Hum Genet* 2018;103:89–99. doi:10.1016/j.ajhg.2018.06.002.
  - [36] Bulik-Sullivan B, Finucane HK, Anttila V, Gusev A, Day FR, Loh P-R, et al. An atlas of genetic correlations across human diseases and traits. *Nat Genet* 2015;47:1236–1241. doi:10.1038/ng.3406.
  - [37] Yang J, Benyamin B, McEvoy BP, Gordon S, Henders AK, Nyholt DR, et al. Common SNPs explain a large proportion of the heritability for human height. *Nat Genet* 2010;42:565–569. doi:10.1038/ng.608.
  - [38] Kääb S, Crawford DC, Sinner MF, Behr ER, Kannankeril PJ, Wilde AAM, et al. A

- Large Candidate Gene Survey Identifies the *KCNE1* D85N Polymorphism as a Possible Modulator of Drug-Induced Torsades de Pointes. *Circ Cardiovasc Genet* 2012;5:91–99. doi:10.1161/CIRCGENETICS.111.960930.
- [39] Moss AJ, Schwartz PJ, Crampton RS, Tzivoni D, Locati EH, MacCluer J, et al. The long QT syndrome. Prospective longitudinal study of 328 families. *Circulation* 1991;84:1136–1144.
- [40] Talmud PJ, Shah S, Whittall R, Futema M, Howard P, Cooper JA, et al. Use of low-density lipoprotein cholesterol gene score to distinguish patients with polygenic and monogenic familial hypercholesterolaemia: a case-control study. *Lancet (London, England)* 2013;381:1293–1301. doi:10.1016/S0140-6736(12)62127-8.
- [41] Rosenberg MA, Lubitz SA, Lin H, Kosova G, Castro VM, Huang P, et al. Validation of Polygenic Scores for QT Interval in Clinical Populations. *Circ Cardiovasc Genet* 2017;10. doi:10.1161/CIRCGENETICS.117.001724.
- [42] Moss AJ, Zareba W, Kaufman ES, Gartman E, Peterson DR, Benhorin J, et al. Increased risk of arrhythmic events in long-QT syndrome with mutations in the pore region of the human ether-a-go-go-related gene potassium channel. *Circulation* 2002;105:794–799.
- [43] Strauss DG, Vicente J, Johannesen L, Blinova K, Mason JW, Weeke P, et al. Common Genetic Variant Risk Score Is Associated With Drug-Induced QT Prolongation and Torsade de Pointes Risk: A Pilot Study. *Circulation* 2017;135:1300–1310. doi:10.1161/CIRCULATIONAHA.116.023980.
- [44] Bagnall RD, Ingles J, Dinger ME, Cowley MJ, Ross SB, Minoche AE, et al. Whole Genome Sequencing Improves Outcomes of Genetic Testing in Patients With Hypertrophic Cardiomyopathy. *J Am Coll Cardiol* 2018;72:419–429. doi:10.1016/j.jacc.2018.04.078.

- [45] Whiffin N, Walsh R, Govind R, Edwards M, Ahmad M, Zhang X, et al. CardioClassifier: disease- and gene-specific computational decision support for clinical genome interpretation. *Genet Med* 2018;20:1246–1254. doi:10.1038/gim.2017.258.
- [46] Walsh R, Mazzarotto F, Whiffin N, Buchan R, Midwinter W, Wilk A, et al. Quantitative approaches to variant classification increase the yield and precision of genetic testing in Mendelian diseases: the case of hypertrophic cardiomyopathy. *Genome Med* 2019;11:5. doi:10.1186/s13073-019-0616-z.
- [47] Nielsen JB, Thorolfsdottir RB, Fritsche LG, Zhou W, Skov MW, Graham SE, et al. Biobank-driven genomic discovery yields new insight into atrial fibrillation biology. *Nat Genet* 2018;50:1234–1239. doi:10.1038/s41588-018-0171-3.
- [48] Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. *Nat Methods* 2012;9:671–675.
- [49] Postema PG, Wilde AAM. The measurement of the QT interval. *Curr Cardiol Rev* 2014;10:287–294.
- [50] Whiffin N, Minikel E, Walsh R, O'Donnell-Luria AH, Karczewski K, Ing AY, et al. Using high-resolution variant frequencies to empower clinical genome interpretation. *Genet Med* 2017;19:1151–1158. doi:10.1038/gim.2017.26.
- [51] Kapplinger JD, Tester DJ, Salisbury BA, Carr JL, Harris-Kerr C, Pollevick GD, et al. Spectrum and prevalence of mutations from the first 2,500 consecutive unrelated patients referred for the FAMILION long QT syndrome genetic test. *Heart Rhythm* 2009;6:1297–1303. doi:10.1016/j.hrthm.2009.05.021.
- [52] Ware JS, Walsh R, Cunningham F, Birney E, Cook SA. Paralogous annotation of disease-causing variants in long QT syndrome genes. *Hum Mutat* 2012;33:1188–1191. doi:10.1002/humu.22114.
- [53] Kapplinger JD, Tseng AS, Salisbury BA, Tester DJ, Callis TE, Alders M, et al.

Enhancing the Predictive Power of Mutations in the C-Terminus of the KCNQ1-Encoded Kv7.1 Voltage-Gated Potassium Channel. *J Cardiovasc Transl Res* 2015;8:187–197. doi:10.1007/s12265-015-9622-8.

- [54] van Setten J, Brody JA, Jamshidi Y, Swenson BR, Butler AM, Campbell H, et al. PR interval genome-wide association meta-analysis identifies 50 loci associated with atrial and atrioventricular electrical activity. *Nat Commun* 2018;9:2904. doi:10.1038/s41467-018-04766-9.
- [55] Sotoodehnia N, Isaacs A, de Bakker PIW, Dörr M, Newton-Cheh C, Nolte IM, et al. Common variants in 22 loci are associated with QRS duration and cardiac ventricular conduction. *Nat Genet* 2010;42:1068–1076. doi:10.1038/ng.716.
- [56] den Hoed M, Eijgelsheim M, Esko T, Brundel BJJM, Peal DS, Evans DM, et al. Identification of heart rate-associated loci and their effects on cardiac conduction and rhythm disorders. *Nat Genet* 2013;45:621–631. doi:10.1038/ng.2610.
- [57] Ramírez J, Duijvenboden S van, Ntalla I, Mifsud B, Warren HR, Tzanis E, et al. Thirty loci identified for heart rate response to exercise and recovery implicate autonomic nervous system. *Nat Commun* 2018;9:1947. doi:10.1038/s41467-018-04148-1.